

CHROMATIC PROPERTIES OF GENERIC PLANAR CONFIGURATIONS OF POINTS

ROLAND BACHER¹ AND DAVID GARBER²

ABSTRACT. We study the Orchard relation defined in [3] for generic configurations of points in the plane (also called order types). We introduce infinitesimally-close points and analyse the relation of this notion with the Orchard relation.

The second part of the paper deals with monochromatic configurations (for the Orchard relation). We give the complete list of all monochromatic configurations up to 7 points and present some constructions and families of monochromatic configurations.

1. INTRODUCTION

A finite set $\mathcal{P} = \{P_1, \dots, P_n\}$ of n points in the oriented affine plane $\mathbb{R}^2$ is a *generic configuration* if three points in $\mathcal{P}$ are never collinear. Two generic configurations of n points $\mathcal{P}^1$ and $\mathcal{P}^2$ are *isotopic* if they can be joined by a continuous curve of generic configurations.

Two generic configurations $\mathcal{P}^1$ and $\mathcal{P}^2$ are *isomorphic* if there exists a bijection $\varphi : \mathcal{P}^1 \rightarrow \mathcal{P}^2$ such that the two triangles having vertices $P, Q, R \in \mathcal{P}^1$ and $\varphi(P), \varphi(Q), \varphi(R) \in \mathcal{P}^2$ induce either always identical or always opposite orientations for all triplets $\{P, Q, R\}$ of points in $\mathcal{P}^1$. In the former case we call $\mathcal{P}^1$ and $\mathcal{P}^2$ *orientedly isomorphic*. The (oriented) isomorphism classes of all generic configurations are called (*oriented*) *order types* by some authors, especially in Computer Sciences (see [1] and [2]).

Isotopic configurations are of course orientedly isomorphic. We ignore to what extent the converse holds.

A line $L \subset \mathbb{R}^2$ *separates* two points $P, Q \in \mathbb{R}^2 \setminus L$ if P and Q are in different connected components of $\mathbb{R}^2 \setminus L$. Given a generic configuration $\mathcal{P}$, we denote by $n(P, Q)$ the number of separating lines defined by pairs of points in $\mathcal{P} \setminus \{P, Q\}$.

¹Support from the Swiss National Science Foundation is gratefully acknowledged.

²Partially supported by the Chateaubriand postdoctoral fellowship funded by the French government.

Date: February 1, 2008.

Definition 1.1 (Orchard relation). *We set $P \sim Q$ if we have $n(P, Q) \equiv (n - 3) \pmod{2}$ for two distinct points $P, Q \in \mathcal{P}$ of a generic configuration $\mathcal{P}$.*

The following result is then a special case of a more general fact, see [3].

Theorem 1.2 (Orchard Theorem). *The Orchard relation is an equivalence relation on the set of points $\mathcal{P}$ of a generic configuration, which consists of at most two classes.*

The name of this result comes from the fact that it yields a canonical rule to plant trees of two species at prescribed generic locations in an orchard.

We call the induced (generally non-trivial) partition the *Orchard partition* of $\mathcal{P}$. Orchard partitions yield invariants for studying generic configurations of points in the affine plane. Iterative use of the Orchard partition, i.e. dividing the set of points into Orchard classes, and iterating this on each class, produces an invariant which distinguishes many pairs of non-isomorphic generic configurations and which is easy to compute and handle.

The present paper is devoted to the illustration of the planar Orchard Theorem applied to generic configurations of n points in the affine plane. Such configurations arise naturally and many features of them have been considered by other authors. This paper yields another such contribution devoted to Orchard properties.

Configurations with trivial Orchard partition (monochromatic configurations) are especially noteworthy to study since they are the “atoms” of the theory. We will describe a few infinite monochromatic families and constructions involving monochromatic configurations.

We introduce and study also some properties of a notion which we call *infinitesimal-closedness*. This property is useful for understanding some monochromatic families and has also some independent interest.

The paper is organized as follows. Section 2 recalls the proof of the Orchard Theorem in the planar case. We also recall and prove the corresponding flip proposition. Section 3 lists all generic configurations having up to 6 points, together with their Orchard partitions. Section 4 deals with infinitesimally-close points and the effect on the Orchard relation after deleting two such points.

The remaining part of the paper is centered on monochromatic configurations. Section 5 gives the complete list of monochromatic configurations up to 7 points. Section 6 deals with some constructions

preserving monochromatic configurations. Section 7 describes some monochromatic families.

Section 8 uses a data-base of Aichholzer, Aurenhammer and Krasser for collecting some statistics related to the Orchard partitions on the set of all generic configurations having at most 9 points.

2. PROOFS OF THE ORCHARD THEOREM AND OF THE FLIP PROPERTY FOR PLANAR CONFIGURATIONS

In this section we prove the planar version of the Orchard Theorem and the flip property along the lines of [3]. The planar setting allows a slight simplification.

Three points P_i, P_j, P_k of a generic planar configuration $\mathcal{P} = \{P_1, \dots, P_n\}$ define three lines which subdivide the projective plane into four triangles: $\Delta_0, \Delta_i, \Delta_j, \Delta_k$, as illustrated in Figure 1.




FIGURE 1. Division of the projective plane into four triangles $\Delta_0 \cup \Delta_i \cup \Delta_j \cup \Delta_k$

Denote by α_i the number of lines connecting two points in $\mathcal{P} \setminus \{P_i, P_j, P_k\}$ and separating (in the affine plane of course) P_i from both points P_j and P_k . The numbers α_j and α_k are defined analogously.

For $* \in \{0, i, j, k\}$ denote by σ_* the number of points in $\mathcal{P} \setminus \{P_i, P_j, P_k\}$ which are in the interior of the triangle Δ_* .

Lemma 2.1. *We have*

$$n(P_i, P_j) = \alpha_i + \alpha_j + \sigma_0 + \sigma_k$$

$$n(P_j, P_k) = \alpha_j + \alpha_k + \sigma_0 + \sigma_i$$

$$n(P_i, P_k) = \alpha_i + \alpha_k + \sigma_0 + \sigma_j$$

Proof. We prove the first formula. The others follow by symmetry.

For a line L separating P_i from P_j we have two cases: $P_k \notin L$ or $P_k \in L$.

In the first case, $P_k \notin L$, the line L separates either P_i from P_k or P_j from P_k (see e.g. line L_1 in Figure 2). Such a line L yields hence a contribution of 1 to either α_i or α_j .

If $P_k \in L$, then L is defined by the point P_k and by another point Q which belongs either to Δ_0 or to Δ_k (consider line L_2 or L_3 in Figure 2).




FIGURE 2. A few lines in a generic configuration

□

Proof of Theorem 1.2. Reflexivity and symmetry of the Orchard relation are obvious. We have to establish transitivity.

Let P_i, P_j, P_k be three points such that $P_i \sim P_j$ and $P_j \sim P_k$. We have to show that $P_i \sim P_k$.

Since $P_i \sim P_j$ and $P_j \sim P_k$ we have $n(P_i, P_j) \equiv (n-3) \pmod{2}$ and $n(P_j, P_k) \equiv (n-3) \pmod{2}$. Adding the equations

$$n(P_i, P_j) = \alpha_i + \alpha_j + \sigma_0 + \sigma_k \equiv (n-3) \pmod{2}$$

$$n(P_j, P_k) = \alpha_j + \alpha_k + \sigma_0 + \sigma_i \equiv (n-3) \pmod{2}$$

obtained by Lemma 2.1, we get

$$\alpha_i + \alpha_k + \sigma_i + \sigma_k \equiv 0 \pmod{2}.$$

The interiors of the four triangles $\Delta_0, \Delta_i, \Delta_j$ and Δ_k contain all points of $\mathcal{P} \setminus \{P_i, P_j, P_k\}$ by genericity and we have hence

$$\sigma_i + \sigma_j + \sigma_k + \sigma_0 = n - 3 .$$

We get therefore

$$n(P_i, P_k) = \alpha_i + \alpha_k + \sigma_0 + \sigma_j \equiv \alpha_i + \alpha_k + \sigma_i + \sigma_k + (n-3) \equiv (n-3) \pmod{2}$$

which shows that $P_i \sim P_k$.

The fact that the Orchard relation has at most two classes follows from the implication $P_i \not\sim P_j$ and $P_j \not\sim P_k \implies P_i \sim P_k$ (the proof of which is, up to a minor change, as above). $\square$

A *flip* is the most elementary move relating non-isomorphic generic configurations:

Definition 2.2 (Flip). *Two generic configurations $\mathcal{P}^1, \mathcal{P}^2$ of n points are related by a flip if there exists a continuous path of configurations $\mathcal{P}(t)$, $-1 \leq t \leq 1$, such that:*

- (1) $\mathcal{P}(-1) = \mathcal{P}^1$ and $\mathcal{P}(1) = \mathcal{P}^2$.
- (2) $\mathcal{P}(t)$ is generic for all $-1 \leq t \leq 1$ except for $t = 0$. The configuration $\mathcal{P}(0)$ has exactly three aligned points, each crossing transversally at $t = 0$ the line spanned by the two other points.

Geometrically, a flip transforms an almost “flat” triangle formed by three points of $\mathcal{P}$ which are nearly aligned into its “mirror”, see Figure 3 for an illustration. We leave it to the reader to show that every pair of generic configurations with n points can be related by a path involving only isotopies and a finite number of flips.




FIGURE 3. Example of a flip

A flip affects the Orchard relation only locally as shown by the following proposition:

Proposition 2.3 (Flip Proposition). *Let $\mathcal{P} = \mathcal{P}(-1) = \{P_1, \dots, P_n\}$ and $\mathcal{P}' = \mathcal{P}(1) = \{P'_1, \dots, P'_n\}$ be two generic configurations related by a flip involving three points $R, S, T \in \mathcal{P}$ corresponding to $R', S', T' \in \mathcal{P}'$. For any two points P, Q we have:*

(1) *if $P, Q \in \{R, S, T\}$ or $P, Q \in \mathcal{P} \setminus \{R, S, T\}$, then:*

$$P \sim Q \quad \text{if and only if} \quad P' \sim' Q'$$

(2) *if $P \in \{R, S, T\}$ and $Q \in \mathcal{P} \setminus \{R, S, T\}$ then:*

$$P \sim Q \quad \text{if and only if} \quad P' \not\sim' Q'$$

Proof. A point $V(t) \in \mathcal{P}(t) \setminus \{R(t), S(t), T(t)\}$ is never aligned with two other points of $\mathcal{P}(t)$. This shows the equality

$$n(P(1), Q(1)) = n(P(-1), Q(-1))$$

for $P(t), Q(t) \in \mathcal{P}(t) \setminus \{R(t), S(t), T(t)\}$ or $P(t), Q(t) \in \{R(t), S(t), T(t)\}$.

On the other hand, $V(t) \in \{R(t), S(t), T(t)\}$ will be involved transversally in the unique alignment $\{R(0), S(0), T(0)\}$ during the flip. For such a point $V(t)$ and $P(t) \in \mathcal{P}(t) \setminus \{R(t), S(t), T(t)\}$ we have hence

$$n(V(1), P(1)) = n(V(-1), P(-1)) \pm 1$$

which proves the result. $\square$

3. ALL GENERIC CONFIGURATIONS UP TO 6 POINTS

This section contains the complete list (up to unoriented isomorphisms) of all generic configurations having at most 6 points together with the corresponding Orchard partitions.

We represent the two classes by black and white vertices. The choice of the black class is of course arbitrary: In case of classes having different cardinalities, black is used for the class containing more elements. In case of a draw, the choice of the black class is arbitrary.

3.1. Generic configurations with up to 5 points. Figures 4 and 5 present all generic configurations (up to non-oriented isomorphism) having at most 5 points, together with their Orchard partitions. Since all these classes admit an orientation-reversing symmetry, these classes coincide with oriented isomorphism classes.




FIGURE 4. All generic configurations with 3 or 4 points




FIGURE 5. All generic configurations with 5 points

3.2. Generic configurations of 6 points. Figures 6, 7 and 8 contain all non-oriented isomorphism classes of generic configurations with 6 points (sorted by the size of their convex hull), together with their Orchard partitions.




FIGURE 6. Generic configurations of 6 points with a convex hull of size 5 or 6

4. INFINITESIMALLY-CLOSE POINTS

In this section, we study configurations containing a pair P, Q of points with $n(P, Q) = 0$ (no separating lines). Such pairs of points will be useful for constructing monochromatic configurations.

Definition 4.1. *Two points P, Q are infinitesimally-close if $n(P, Q) = 0$.*

Remark 4.2. *Infinitesimally-close points are Orchard-equivalent if the total number of points in the configuration is odd and they are not equivalent otherwise.*

The following proposition describes the change in the Orchard relation induced by deleting two infinitesimally-close points:

Proposition 4.3. *Let $R, S \in \mathcal{P}$ be two infinitesimally-close points of a generic configuration $\mathcal{P}$ having n points. Denote by L the line containing R and S and by $\mathcal{P}' = \mathcal{P} \setminus \{R, S\}$ the subconfiguration of $\mathcal{P}$ after deletion of R and S .*

- (1) *If $P, Q \in \mathcal{P} \setminus \{R, S\}$ are not separated by L , then*

$$P \sim_{\mathcal{P}} Q \quad \text{if and only if} \quad P \sim_{\mathcal{P}'} Q .$$

- (2) *If $P, Q \in \mathcal{P} \setminus \{R, S\}$ are separated by L , then*

$$P \sim_{\mathcal{P}} Q \quad \text{if and only if} \quad P \not\sim_{\mathcal{P}'} Q .$$




FIGURE 7. Generic configurations of 6 points with a convex hull of size 4

Proof. If P, Q are not separated by L , then $n_{\mathcal{P}}(P, Q)$ and $n_{\mathcal{P}'}(P, Q)$ have the same parity. Indeed, if a separating line between P and Q is generated by R (say) and a second point $T \in \mathcal{P} \setminus \{R, S, P, Q\}$, then the line generated by S and T separates P and Q too. The number of such lines is hence even.




FIGURE 8. Generic configurations of 6 points with a convex hull of size 3

If the line L spanned by R and S separates two points $P, Q \in \mathcal{P} \setminus \{R, S\}$ then the number $n_{\mathcal{P}}(P, Q) - n_{\mathcal{P}'}(P, Q)$ is odd since it contains, together with all pairs of lines mentioned above, also the line L . $\square$

Remark 4.4. *Generic configurations consisting of few points have generally pairs of infinitesimally-close points. Generic configurations having a huge number of points however have only rarely infinitesimally-close points.*

4.1. Infinitesimal contractions and infinitesimally equivalent configurations. Let $\mathcal{P}$ be a configuration with two infinitesimally-close points R and S . This implies that the two configurations $\mathcal{P} \setminus \{R\}$ and $\mathcal{P} \setminus \{S\}$, obtained by deleting either R or S from $\mathcal{P}$, are isotopic. We call the isotopy class $\mathcal{P}'$ of $\mathcal{P} \setminus \{R\}$ the *infinitesimal contraction* of $\mathcal{P}$ along $[R, S]$. Two configurations $\mathcal{P}$, $\mathcal{P}'$ related through an infinitesimal contraction are *infinitesimally related*. Two configurations $\mathcal{P}^1$ and $\mathcal{P}^2$ which can be joined by a path of infinitesimally related configurations are *infinitesimally equivalent* and the set of all configurations infinitesimally equivalent to $\mathcal{P}$ forms of course an equivalence class. Nothing interesting can be said concerning Orchard properties of such a class since removal of a point changes the Orchard relation dramatically. It has however a nice property given by the following result.

Theorem 4.5. *Each equivalence class of infinitesimally related configurations contains a unique (up to isomorphism) minimal representative. The minimal representative of a configuration $\mathcal{P}$ can always be constructed as a subconfiguration reached through a succession of infinitesimal contractions starting at $\mathcal{P}$.*

Proof. We have to prove unicity of a minimal representative. Choose a minimal sequence $\gamma(0), \gamma(1), \dots, \gamma(l)$ with $\gamma(i)$ minimally adjacent to $\gamma(i+1)$ relating two such minimal representatives. By minimality of $\gamma(0)$ and $\gamma(l)$ there is a first index i such that $\gamma(i+1)$ is an infinitesimal contraction of $\gamma(i)$ along $[R, S]$. This implies that the point R or S has been added before, thus contradicting the minimality of the sequence $\gamma(0), \dots, \gamma(l)$. $\square$

Remark 4.6. (1) *Infinitesimal contractions admit a sort of inverse: Given a generic configuration $\mathcal{P}$ of n points, choose a point $R \in \mathcal{P}$ and one of the $(n-1)$ opposite pairs of cones delimited by lines through R and $\mathcal{P} \setminus \{R\}$. Add now a new point S very close to R in the interior of one of the two chosen opposite cones. This clearly yields a configuration with $n(R, S) = 0$ and every configuration of $(n+1)$ points infinitesimally adjacent to $\mathcal{P}$ can be constructed in this way.*

(2) *The notion of infinitesimal equivalence behaves well with respect to isotopy: If two generic configurations are isomorphic but not*

isotopic, then their minimal representatives are also only isotopic but not isotopic.

5. MONOCHROMATIC CONFIGURATIONS

This section is devoted to the study of *monochromatic configurations* which are configurations with trivial Orchard partition.

Figures 9-12 contain all monochromatic configurations (up to non-oriented isomorphism) having at most 7 points.




FIGURE 9. Monochromatic configurations with up to 6 points




FIGURE 10. Monochromatic configurations of 7 points having convex hull of size 5, 6 or 7




FIGURE 11. Monochromatic configurations of 7 points with convex hull of size 4




FIGURE 12. Monochromatic configurations of 7 points with convex hull of size 3

6. MONOCHROMATIC CONSTRUCTIONS

This section contains a few constructions of monochromatic configurations.

Proposition 6.1. *Let $\mathcal{P}$ be a monochromatic configuration having two infinitesimally-close points P and Q in its convex hull. Then $\mathcal{P} \setminus \{P, Q\}$ is also monochromatic.*

Proof. Use Proposition 4.3. □

This proposition can be used as follows:

Construction 6.2. *Let $\mathcal{P} = \mathcal{P}' \cup \{P, Q\}$ be a generic configuration with an odd number of points having two infinitesimally-close vertices P, Q in its convex hull. Suppose that $\mathcal{P}'$ is monochromatic and $n(P, P_0)$ is even for some point $P_0 \in \mathcal{P}'$. Then $\mathcal{P}$ is monochromatic.*

Proof. Apply the previous proposition. □

Remark 6.3. *The same construction works also for a configuration $\mathcal{P}$ with an Orchard partition (n, n) : Add (generically) two infinitesimally-close points P, Q to $\mathcal{P}$ such that P, Q are in the convex hull of $\mathcal{P} \cup \{P, Q\}$. The configuration $\mathcal{P} \cup \{P, Q\}$ has then Orchard partition $(n + 1, n + 1)$.*

There are of course many more constructions. Let us close this section by mentioning another one:

Given a monochromatic configuration $\mathcal{P}$ having an odd number of points, add generically two infinitesimally-close points P, Q . This yields a configuration $\mathcal{P}'$ having two classes $\mathcal{C}^1$ and $\mathcal{C}^2$, with $P, Q \in \mathcal{C}^1$, which are separated by the line L through P, Q . Add now another pair of infinitesimally-close points P', Q' defining a line L' parallel to L in such a way that L' separates P, Q from all points in $\mathcal{C}^2$. In the resulting configuration, the classes $\mathcal{C}^1$ and $\mathcal{C}^2$ merge. Only the new points P', Q' might be outside this class. In this case, shift them along L' into a suitable position.

7. SOME FAMILIES OF MONOCHROMATIC CONFIGURATIONS

In this section we describe some monochromatic families.

Proposition 7.1. *Vertices of a convex polygon having $2n + 1$ edges yield a monochromatic configuration.*

Proof. Two adjacent vertices P and Q of the polygon satisfy clearly $n(P, Q) = 0$. This implies $P \sim Q$ since n is odd. $\square$

Remark 7.2. *The above family can also be obtained by iterating Construction 6.2 starting with the unique configuration consisting of one point.*

Remark 7.3. *The same argument shows that vertices of a convex polygon with $2n$ edges have alternating colours.*

Another monochromatic family can be described as follows:

Proposition 7.4. *The following configuration $\mathcal{P}$ of $2n$ points is monochromatic for $n \geq 3$: Consider n vertices spanning a regular polygon R (any convex polygon will in fact work) having n sides. Add n more points which are interior to R and very close to the midpoints of all edges in R (Figure 13 displays the example $n = 5$).*

Proof. Consider first two adjacent vertices P and Q of the polygon R . Denote by $S \in \mathcal{P}$ the point near the barycenter $(P + Q)/2$ of P and Q . The points P and Q are separated by all lines through S and one of the remaining $2n - 3$ points of $\mathcal{P} \setminus \{P, Q, S\}$. Moreover, there are two more separating lines going through vertices adjacent to P and Q and through the corresponding near-midpoints of the edges adjacent to the edge $[P, Q]$. We have hence $n(P, Q) = 2n - 1 \equiv 2n - 3 \pmod{2}$ which shows that $P \sim Q$.

The points P and S are separated by exactly one of the two extra lines mentioned above and hence we have $P \sim S$ too. $\square$




FIGURE 13. An example for a member in the family for $n = 5$

Remark 7.5. One can prove the last result also by starting with a convex polygon having $2n$ vertices $\{s_0, \dots, s_{2n-1}\}$. By Remark 7.3, we have $s_a \equiv s_b$ if and only if $a \equiv b \pmod{2}$. Perform now flips with respect to all triangles of vertices $s_{2i}, s_{2i+1}, s_{2i+2}$, $0 \leq i \leq n-1$ (with indices read $\pmod{2n}$). A vertex s_k is hence involved in either 2 or 1 flips according to the parity of k and the resulting final configuration is monochromatic by the Flip Proposition (Proposition 2.3).

A monochromatic family involving congruences is given as follows:

Proposition 7.6. Let $n \geq 5$ be an odd number, such that $(n+1) \equiv 0 \pmod{8}$ or $(n+1) \equiv 6 \pmod{8}$. The n vertices of a regular polygon having n sides together with its barycenter yield a monochromatic configuration $\mathcal{P}$ of $(n+1)$ points (Figure 14 shows the case $n = 5$).




FIGURE 14. An example for a member in the family for $n = 5$

Proof. It is easy to check that $n(P, Q) = 1$ for two adjacent vertices of the regular n -gon.

We have yet to show that $n(P, C)$ is odd when P is a vertex of the regular n -gon and C is its barycenter. Any line L separating C from

P comes together with its mirror obtained by reflecting L with respect to the line joining C and P (see lines L_1 and L_2 in Figure 15). Only separating lines orthogonal to the segment $[C, P]$ (like for instance line L_3 in Figure 15) have hence to be taken into account when counting separating lines $\pmod{2}$.




FIGURE 15. Examples for some separating lines for the segment $[P_1, C]$, where $n = 7$

Such lines are in bijection with the set

$$\{k \in \mathbb{N} \mid 0 < k \frac{2\pi}{n} < \frac{\pi}{2}\} = \{1, 2, \dots, \lfloor \frac{n}{4} \rfloor\},$$

and the result follows. $\square$

Remark 7.7. For $n \equiv 1 \pmod{8}$ or $n \equiv 3 \pmod{8}$, the configurations considered in Proposition 7.6 have Orchard partition $(n, 1)$.

A somewhat similar family is given by:

Proposition 7.8. Let $n \geq 3$ be an odd number. The following construction yields a monochromatic configuration of $2n + 1$ points: Draw the n vertices $\tilde{P}_1, \dots, \tilde{P}_n$ of a regular n -gon. Now replace each vertex $\tilde{P}_i$ by two infinitesimally-close points P_i, P'_i , in order to obtain a configuration with convex hull of size $2n$. Add now a last point C at the barycenter of this $2n$ -gon (See Figure 10 number (2) for an example for $n = 3$).

Proof. We have to check that $n(P_i, P'_i)$, $n(P'_i, P_{i+1})$ and $n(P_1, C)$ are even.

It is obvious that $n(P_i, P'_i) = 0$. Moreover, $n(P'_i, P_{i+1}) = 2$ with two separating lines defined by the midpoint C and the two vertices opposite to the segment $[P'_i, P_{i+1}]$.

It remains to check that $n(C, P_i)$ is even. This follows from the fact that each line separating C from P_i has at least one of its defining points in a pair of infinitesimally-close points different from P_i and P'_i . $\square$

8. SOME STATISTICS

Aichholzer, Aurenhammer and Krasser have computed a data-base of all generic configurations having $n \leq 10$ points, up to unoriented isomorphism (see [1] and [2]). Assuming completeness of their data-base, we have computed the Orchard partitions for all generic configurations having $n \leq 9$ points. The results for configurations of 7, 8 and 9 points are presented in Tables 1-3 according to the size of the configuration's convex hull and the size of the two equivalence classes of the Orchard partition.

Monochromatic configurations are of course partitions of type $(0, *)$.

Partition:	(0,7)	(1,6)	(2,5)	(3,4)
Convex hull	mono			
3	3	7	13	26
4	4	11	16	28
5	1	3	5	13
6	1	0	1	2
7	1	0	0	0
Sum:	10	21	35	69

TABLE 1. Statistics for configurations with 7 points

Partition:	(0,8)	(1,7)	(2,6)	(3,5)	(4,4)
Convex hull	mono				
3	10	38	252	552	326
4	12	92	323	635	406
5	4	29	87	261	189
6	1	4	11	38	36
7	1	1	0	3	3
8	0	0	0	0	1
Sum:	28	164	673	1489	961

TABLE 2. Statistics for configurations with 8 points

Partition: Convex hull	(0,9) mono	(1,8)	(2,7)	(3,6)	(4,5)
3	272	2469	8459	17493	26542
4	306	2484	10012	23234	34439
5	231	1277	4184	9273	13267
6	52	230	661	1490	2119
7	6	13	42	102	148
8	1	0	3	3	4
9	1	0	0	0	0
Sum:	869	6473	23361	51595	76519

TABLE 3. Statistics for configurations with 9 points

ACKNOWLEDGMENT

The second author wishes to thank Mikhail Zaidenberg and the Institut Fourier for hosting his stay.

REFERENCES

- [1] Aichholzer, O., Aurenhammer, F. and Krasser, H., *Enumerating order types for small point sets with applications*, in: Proc. 17th Ann. ACM Symp. Computational Geometry, 11–18, Medford, Massachusetts, USA, 2001.
- [2] Aichholzer, O. and Krasser, H., *The point set order type data base: A collection of applications and results*, in: Proc. 13th Annual Canadian Conference on Computational Geometry CCCG 2001, 17–20, Waterloo, Ontario, Canada, 2001.
- [3] Bacher, R., *An Orchard Theorem*, preprint [math.CO/0206266].

INSTITUT FOURIER, BP 74, 38402 SAINT-MARTIN D'HERES CEDEX, FRANCE
E-mail address: {bacher,garber}@mozart.ujf-grenoble.fr